




United States
Department of
Agriculture

NRCS

Natural
Resources
Conservation
Service

A product of the National
Cooperative Soil Survey,
a joint effort of the United
States Department of
Agriculture and other
Federal agencies, State
agencies including the
Agricultural Experiment
Stations, and local
participants

Custom Soil Resource Report for **Jackson County, Texas**



Preface

Soil surveys contain information that affects land use planning in survey areas. They highlight soil limitations that affect various land uses and provide information about the properties of the soils in the survey areas. Soil surveys are designed for many different users, including farmers, ranchers, foresters, agronomists, urban planners, community officials, engineers, developers, builders, and home buyers. Also, conservationists, teachers, students, and specialists in recreation, waste disposal, and pollution control can use the surveys to help them understand, protect, or enhance the environment.

Various land use regulations of Federal, State, and local governments may impose special restrictions on land use or land treatment. Soil surveys identify soil properties that are used in making various land use or land treatment decisions. The information is intended to help the land users identify and reduce the effects of soil limitations on various land uses. The landowner or user is responsible for identifying and complying with existing laws and regulations.

Although soil survey information can be used for general farm, local, and wider area planning, onsite investigation is needed to supplement this information in some cases. Examples include soil quality assessments (<http://www.nrcs.usda.gov/wps/portal/nrcs/main/soils/health/>) and certain conservation and engineering applications. For more detailed information, contact your local USDA Service Center (<https://offices.sc.egov.usda.gov/locator/app?agency=nrcs>) or your NRCS State Soil Scientist (http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/contactus/?cid=nrcs142p2_053951).

Great differences in soil properties can occur within short distances. Some soils are seasonally wet or subject to flooding. Some are too unstable to be used as a foundation for buildings or roads. Clayey or wet soils are poorly suited to use as septic tank absorption fields. A high water table makes a soil poorly suited to basements or underground installations.

The National Cooperative Soil Survey is a joint effort of the United States Department of Agriculture and other Federal agencies, State agencies including the Agricultural Experiment Stations, and local agencies. The Natural Resources Conservation Service (NRCS) has leadership for the Federal part of the National Cooperative Soil Survey.

Information about soils is updated periodically. Updated information is available through the NRCS Web Soil Survey, the site for official soil survey information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or a part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require

alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410 or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.

Contents

Preface	2
How Soil Surveys Are Made	5
Soil Map	8
Soil Map.....	9
Legend.....	10
Map Unit Legend.....	11
Map Unit Descriptions.....	11
Jackson County, Texas.....	13
DaA—Dacosta sandy clay loam, 0 to 1 percent slopes.....	13
InB—Inez fine sandy loam, 0 to 2 percent slopes.....	14
KuC—Kuy sand, 1 to 5 percent slopes.....	16
LaA—Laewest clay, 0 to 1 percent slopes.....	17
MaC—Marcado sandy clay loam, 3 to 8 percent slopes.....	19
MbB—Milby sand, 0 to 2 percent slopes.....	20
MrA—Morales-Cieno complex, 0 to 1 percent slopes, frequently ponded..	21
NcA—Nada-Cieno complex, 0 to 1 percent slopes, frequently ponded.....	24
Nv—Navidad fine sandy loam, frequently flooded.....	26

How Soil Surveys Are Made

Soil surveys are made to provide information about the soils and miscellaneous areas in a specific area. They include a description of the soils and miscellaneous areas and their location on the landscape and tables that show soil properties and limitations affecting various uses. Soil scientists observed the steepness, length, and shape of the slopes; the general pattern of drainage; the kinds of crops and native plants; and the kinds of bedrock. They observed and described many soil profiles. A soil profile is the sequence of natural layers, or horizons, in a soil. The profile extends from the surface down into the unconsolidated material in which the soil formed or from the surface down to bedrock. The unconsolidated material is devoid of roots and other living organisms and has not been changed by other biological activity.

Currently, soils are mapped according to the boundaries of major land resource areas (MLRAs). MLRAs are geographically associated land resource units that share common characteristics related to physiography, geology, climate, water resources, soils, biological resources, and land uses (USDA, 2006). Soil survey areas typically consist of parts of one or more MLRA.

The soils and miscellaneous areas in a survey area occur in an orderly pattern that is related to the geology, landforms, relief, climate, and natural vegetation of the area. Each kind of soil and miscellaneous area is associated with a particular kind of landform or with a segment of the landform. By observing the soils and miscellaneous areas in the survey area and relating their position to specific segments of the landform, a soil scientist develops a concept, or model, of how they were formed. Thus, during mapping, this model enables the soil scientist to predict with a considerable degree of accuracy the kind of soil or miscellaneous area at a specific location on the landscape.

Commonly, individual soils on the landscape merge into one another as their characteristics gradually change. To construct an accurate soil map, however, soil scientists must determine the boundaries between the soils. They can observe only a limited number of soil profiles. Nevertheless, these observations, supplemented by an understanding of the soil-vegetation-landscape relationship, are sufficient to verify predictions of the kinds of soil in an area and to determine the boundaries.

Soil scientists recorded the characteristics of the soil profiles that they studied. They noted soil color, texture, size and shape of soil aggregates, kind and amount of rock fragments, distribution of plant roots, reaction, and other features that enable them to identify soils. After describing the soils in the survey area and determining their properties, the soil scientists assigned the soils to taxonomic classes (units). Taxonomic classes are concepts. Each taxonomic class has a set of soil characteristics with precisely defined limits. The classes are used as a basis for comparison to classify soils systematically. Soil taxonomy, the system of taxonomic classification used in the United States, is based mainly on the kind and character of soil properties and the arrangement of horizons within the profile. After the soil

Custom Soil Resource Report

scientists classified and named the soils in the survey area, they compared the individual soils with similar soils in the same taxonomic class in other areas so that they could confirm data and assemble additional data based on experience and research.

The objective of soil mapping is not to delineate pure map unit components; the objective is to separate the landscape into landforms or landform segments that have similar use and management requirements. Each map unit is defined by a unique combination of soil components and/or miscellaneous areas in predictable proportions. Some components may be highly contrasting to the other components of the map unit. The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The delineation of such landforms and landform segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, onsite investigation is needed to define and locate the soils and miscellaneous areas.

Soil scientists make many field observations in the process of producing a soil map. The frequency of observation is dependent upon several factors, including scale of mapping, intensity of mapping, design of map units, complexity of the landscape, and experience of the soil scientist. Observations are made to test and refine the soil-landscape model and predictions and to verify the classification of the soils at specific locations. Once the soil-landscape model is refined, a significantly smaller number of measurements of individual soil properties are made and recorded. These measurements may include field measurements, such as those for color, depth to bedrock, and texture, and laboratory measurements, such as those for content of sand, silt, clay, salt, and other components. Properties of each soil typically vary from one point to another across the landscape.

Observations for map unit components are aggregated to develop ranges of characteristics for the components. The aggregated values are presented. Direct measurements do not exist for every property presented for every map unit component. Values for some properties are estimated from combinations of other properties.

While a soil survey is in progress, samples of some of the soils in the area generally are collected for laboratory analyses and for engineering tests. Soil scientists interpret the data from these analyses and tests as well as the field-observed characteristics and the soil properties to determine the expected behavior of the soils under different uses. Interpretations for all of the soils are field tested through observation of the soils in different uses and under different levels of management. Some interpretations are modified to fit local conditions, and some new interpretations are developed to meet local needs. Data are assembled from other sources, such as research information, production records, and field experience of specialists. For example, data on crop yields under defined levels of management are assembled from farm records and from field or plot experiments on the same kinds of soil.

Predictions about soil behavior are based not only on soil properties but also on such variables as climate and biological activity. Soil conditions are predictable over long periods of time, but they are not predictable from year to year. For example, soil scientists can predict with a fairly high degree of accuracy that a given soil will have a high water table within certain depths in most years, but they cannot predict that a high water table will always be at a specific level in the soil on a specific date.

After soil scientists located and identified the significant natural bodies of soil in the survey area, they drew the boundaries of these bodies on aerial photographs and


Custom Soil Resource Report

identified each as a specific map unit. Aerial photographs show trees, buildings, fields, roads, and rivers, all of which help in locating boundaries accurately.


Soil Map

The soil map section includes the soil map for the defined area of interest, a list of soil map units on the map and extent of each map unit, and cartographic symbols displayed on the map. Also presented are various metadata about data used to produce the map, and a description of each soil map unit.

Custom Soil Resource Report Soil Map



Map Scale: 1:44,700 if printed on A landscape (11" x 8.5") sheet.



Map projection: Web Mercator Corner coordinates: WGS84 Edge tics: UTM Zone 14N WGS84

MAP LEGEND

Area of Interest (AOI)

 Area of Interest (AOI)

Soils

 Soil Map Unit Polygons

 Soil Map Unit Lines

 Soil Map Unit Points

Special Point Features

-  Blowout
-  Borrow Pit
-  Clay Spot
-  Closed Depression
-  Gravel Pit
-  Gravelly Spot
-  Landfill
-  Lava Flow
-  Marsh or swamp
-  Mine or Quarry
-  Miscellaneous Water
-  Perennial Water
-  Rock Outcrop
-  Saline Spot
-  Sandy Spot
-  Severely Eroded Spot
-  Sinkhole
-  Slide or Slip
-  Sodic Spot

-  Spoil Area
-  Stony Spot
-  Very Stony Spot
-  Wet Spot
-  Other
-  Special Line Features

Water Features

 Streams and Canals

Transportation

-  Rails
-  Interstate Highways
-  US Routes
-  Major Roads
-  Local Roads

Background

 Aerial Photography

MAP INFORMATION

The soil surveys that comprise your AOI were mapped at 1:24,000.

Please rely on the bar scale on each map sheet for map measurements.

Source of Map: Natural Resources Conservation Service
 Web Soil Survey URL:
 Coordinate System: Web Mercator (EPSG:3857)

Maps from the Web Soil Survey are based on the Web Mercator projection, which preserves direction and shape but distorts distance and area. A projection that preserves area, such as the Albers equal-area conic projection, should be used if more accurate calculations of distance or area are required.

This product is generated from the USDA-NRCS certified data as of the version date(s) listed below.

Soil Survey Area: Jackson County, Texas
 Survey Area Data: Version 12, Sep 21, 2016

Soil map units are labeled (as space allows) for map scales 1:50,000 or larger.

Date(s) aerial images were photographed: Jan 27, 2011—May 14, 2011

The orthophoto or other base map on which the soil lines were compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident.

Map Unit Legend

Jackson County, Texas (TX239)			
Map Unit Symbol	Map Unit Name	Acres in AOI	Percent of AOI
DaA	Dacosta sandy clay loam, 0 to 1 percent slopes	764.3	14.7%
InB	Inez fine sandy loam, 0 to 2 percent slopes	1,237.4	23.8%
KuC	Kuy sand, 1 to 5 percent slopes	15.7	0.3%
LaA	Laewest clay, 0 to 1 percent slopes	111.6	2.1%
MaC	Marcado sandy clay loam, 3 to 8 percent slopes	416.5	8.0%
MbB	Milby sand, 0 to 2 percent slopes	194.1	3.7%
MrA	Morales-Cieno complex, 0 to 1 percent slopes, frequently ponded	933.8	17.9%
NcA	Nada-Cieno complex, 0 to 1 percent slopes, frequently ponded	1,224.1	23.5%
Nv	Navidad fine sandy loam, frequently flooded	306.0	5.9%
Totals for Area of Interest		5,203.6	100.0%

Map Unit Descriptions

The map units delineated on the detailed soil maps in a soil survey represent the soils or miscellaneous areas in the survey area. The map unit descriptions, along with the maps, can be used to determine the composition and properties of a unit.

A map unit delineation on a soil map represents an area dominated by one or more major kinds of soil or miscellaneous areas. A map unit is identified and named according to the taxonomic classification of the dominant soils. Within a taxonomic class there are precisely defined limits for the properties of the soils. On the landscape, however, the soils are natural phenomena, and they have the characteristic variability of all natural phenomena. Thus, the range of some observed properties may extend beyond the limits defined for a taxonomic class. Areas of soils of a single taxonomic class rarely, if ever, can be mapped without including areas of other taxonomic classes. Consequently, every map unit is made up of the soils or miscellaneous areas for which it is named and some minor components that belong to taxonomic classes other than those of the major soils.

Most minor soils have properties similar to those of the dominant soil or soils in the map unit, and thus they do not affect use and management. These are called noncontrasting, or similar, components. They may or may not be mentioned in a particular map unit description. Other minor components, however, have properties and behavioral characteristics divergent enough to affect use or to require different

Custom Soil Resource Report

management. These are called contrasting, or dissimilar, components. They generally are in small areas and could not be mapped separately because of the scale used. Some small areas of strongly contrasting soils or miscellaneous areas are identified by a special symbol on the maps. If included in the database for a given area, the contrasting minor components are identified in the map unit descriptions along with some characteristics of each. A few areas of minor components may not have been observed, and consequently they are not mentioned in the descriptions, especially where the pattern was so complex that it was impractical to make enough observations to identify all the soils and miscellaneous areas on the landscape.

The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The objective of mapping is not to delineate pure taxonomic classes but rather to separate the landscape into landforms or landform segments that have similar use and management requirements. The delineation of such segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, however, onsite investigation is needed to define and locate the soils and miscellaneous areas.

An identifying symbol precedes the map unit name in the map unit descriptions. Each description includes general facts about the unit and gives important soil properties and qualities.

Soils that have profiles that are almost alike make up a *soil series*. Except for differences in texture of the surface layer, all the soils of a series have major horizons that are similar in composition, thickness, and arrangement.

Soils of one series can differ in texture of the surface layer, slope, stoniness, salinity, degree of erosion, and other characteristics that affect their use. On the basis of such differences, a soil series is divided into *soil phases*. Most of the areas shown on the detailed soil maps are phases of soil series. The name of a soil phase commonly indicates a feature that affects use or management. For example, Alpha silt loam, 0 to 2 percent slopes, is a phase of the Alpha series.

Some map units are made up of two or more major soils or miscellaneous areas. These map units are complexes, associations, or undifferentiated groups.

A *complex* consists of two or more soils or miscellaneous areas in such an intricate pattern or in such small areas that they cannot be shown separately on the maps. The pattern and proportion of the soils or miscellaneous areas are somewhat similar in all areas. Alpha-Beta complex, 0 to 6 percent slopes, is an example.

An *association* is made up of two or more geographically associated soils or miscellaneous areas that are shown as one unit on the maps. Because of present or anticipated uses of the map units in the survey area, it was not considered practical or necessary to map the soils or miscellaneous areas separately. The pattern and relative proportion of the soils or miscellaneous areas are somewhat similar. Alpha-Beta association, 0 to 2 percent slopes, is an example.

An *undifferentiated group* is made up of two or more soils or miscellaneous areas that could be mapped individually but are mapped as one unit because similar interpretations can be made for use and management. The pattern and proportion of the soils or miscellaneous areas in a mapped area are not uniform. An area can be made up of only one of the major soils or miscellaneous areas, or it can be made up of all of them. Alpha and Beta soils, 0 to 2 percent slopes, is an example.

Some surveys include *miscellaneous areas*. Such areas have little or no soil material and support little or no vegetation. Rock outcrop is an example.

Jackson County, Texas

DaA—Dacosta sandy clay loam, 0 to 1 percent slopes

Map Unit Setting

National map unit symbol: 2v3c1
Elevation: 10 to 220 feet
Mean annual precipitation: 30 to 49 inches
Mean annual air temperature: 69 to 71 degrees F
Frost-free period: 270 to 300 days
Farmland classification: All areas are prime farmland

Map Unit Composition

Dacosta and similar soils: 90 percent
Minor components: 10 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Dacosta

Setting

Landform: Flats
Landform position (three-dimensional): Talf
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Clayey fluviomarine deposits derived from igneous, metamorphic and sedimentary rock

Typical profile

A - 0 to 10 inches: sandy clay loam
Bt1 - 10 to 22 inches: sandy clay loam
Bt2 - 22 to 43 inches: clay
Btk - 43 to 74 inches: sandy clay
2BCt - 74 to 80 inches: sandy clay loam

Properties and qualities

Slope: 0 to 1 percent
Depth to restrictive feature: More than 80 inches
Natural drainage class: Moderately well drained
Runoff class: Medium
Capacity of the most limiting layer to transmit water (Ksat): Very low to moderately low (0.00 to 0.06 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Calcium carbonate, maximum in profile: 5 percent
Salinity, maximum in profile: Very slightly saline to moderately saline (2.0 to 8.0 mmhos/cm)
Sodium adsorption ratio, maximum in profile: 11.0
Available water storage in profile: Moderate (about 6.8 inches)

Interpretive groups

Land capability classification (irrigated): 3s
Land capability classification (nonirrigated): 3s
Hydrologic Soil Group: D
Ecological site: Blackland 24-44" PZ (R150AY526TX)

Custom Soil Resource Report

Hydric soil rating: No

Minor Components

Laewest

Percent of map unit: 5 percent
Landform: Flats
Landform position (three-dimensional): Talf
Microfeatures of landform position: Gilgai
Down-slope shape: Linear
Across-slope shape: Linear
Ecological site: Blackland 24-44" PZ (R150AY526TX)
Hydric soil rating: No

Edna

Percent of map unit: 3 percent
Landform: Flats
Landform position (three-dimensional): Talf
Down-slope shape: Linear
Across-slope shape: Linear
Ecological site: Tight Sandy Loam 25-35" PZ (R150AY646TX)
Hydric soil rating: No

Cieno

Percent of map unit: 1 percent
Landform: — error in exists on —
Landform position (three-dimensional): Dip
Down-slope shape: Concave
Across-slope shape: Concave
Ecological site: Lowland 35-56" PZ (R150AY537TX)
Hydric soil rating: Yes

Telferner

Percent of map unit: 1 percent
Landform: Ridges
Landform position (three-dimensional): Rise
Down-slope shape: Linear
Across-slope shape: Convex
Ecological site: Loamy Prairie 28-44 PZ (R150AY535TX)
Hydric soil rating: No

InB—Inez fine sandy loam, 0 to 2 percent slopes

Map Unit Setting

National map unit symbol: 2vv31
Elevation: 10 to 400 feet
Mean annual precipitation: 32 to 44 inches
Mean annual air temperature: 68 to 72 degrees F
Frost-free period: 260 to 300 days
Farmland classification: Farmland of statewide importance

Map Unit Composition

Inez and similar soils: 85 percent

Minor components: 15 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Inez

Setting

Landform: Stream terraces

Landform position (three-dimensional): Tread

Down-slope shape: Linear

Across-slope shape: Linear

Parent material: Late pleistocene loamy alluvium

Typical profile

A - 0 to 8 inches: fine sandy loam

E - 8 to 14 inches: fine sandy loam

Bt - 14 to 42 inches: clay

Btk - 42 to 80 inches: clay loam

Properties and qualities

Slope: 0 to 2 percent

Depth to restrictive feature: More than 80 inches

Natural drainage class: Moderately well drained

Runoff class: Very high

Capacity of the most limiting layer to transmit water (Ksat): Very low to moderately low (0.00 to 0.06 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Calcium carbonate, maximum in profile: 5 percent

Salinity, maximum in profile: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)

Sodium adsorption ratio, maximum in profile: 6.0

Available water storage in profile: Moderate (about 8.8 inches)

Interpretive groups

Land capability classification (irrigated): 3s

Land capability classification (nonirrigated): 3s

Hydrologic Soil Group: D

Ecological site: Sandy Loam 28-44" PZ (R150AY542TX)

Hydric soil rating: No

Minor Components

Cieno

Percent of map unit: 4 percent

Landform: — error in exists on —

Landform position (three-dimensional): Dip

Down-slope shape: Concave

Across-slope shape: Concave

Ecological site: Lowland 35-56" PZ (R150AY537TX)

Hydric soil rating: Yes

Milby

Percent of map unit: 4 percent

Custom Soil Resource Report

Landform: Terraces
Landform position (three-dimensional): Tread
Down-slope shape: Linear
Across-slope shape: Convex
Ecological site: SANDY 25-35" PZ (R150AY644TX)
Hydric soil rating: No

Edna

Percent of map unit: 4 percent
Landform: Flats
Landform position (three-dimensional): Talf
Down-slope shape: Linear
Across-slope shape: Linear
Ecological site: Tight Sandy Loam 25-35" PZ (R150AY646TX)
Hydric soil rating: No

Fordtran

Percent of map unit: 3 percent
Landform: Terraces
Landform position (three-dimensional): Tread
Down-slope shape: Linear
Across-slope shape: Convex
Ecological site: Sandy Prairie 25-35" PZ (R150AY543TX)
Hydric soil rating: No

KuC—Kuy sand, 1 to 5 percent slopes

Map Unit Setting

National map unit symbol: dlp0
Elevation: 80 to 300 feet
Mean annual precipitation: 36 to 42 inches
Mean annual air temperature: 70 to 72 degrees F
Frost-free period: 260 to 300 days
Farmland classification: Not prime farmland

Map Unit Composition

Kuy and similar soils: 80 percent
Minor components: 20 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Kuy

Setting

Landform: Terraces
Landform position (three-dimensional): Tread
Down-slope shape: Convex
Across-slope shape: Convex
Parent material: Loamy and sandy alluvium of pleistocene age

Typical profile

H1 - 0 to 52 inches: sand

Custom Soil Resource Report

H2 - 52 to 86 inches: clay loam

Properties and qualities

Slope: 1 to 5 percent

Depth to restrictive feature: More than 80 inches

Natural drainage class: Moderately well drained

Runoff class: Negligible

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 1.98 in/hr)

Depth to water table: About 48 to 72 inches

Frequency of flooding: None

Frequency of ponding: None

Salinity, maximum in profile: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)

Sodium adsorption ratio, maximum in profile: 2.0

Available water storage in profile: Low (about 5.9 inches)

Interpretive groups

Land capability classification (irrigated): 3e

Land capability classification (nonirrigated): 3e

Hydrologic Soil Group: A

Ecological site: Deep Sand 35-42" PZ (R150AY532TX)

Hydric soil rating: No

Minor Components

Milby

Percent of map unit: 8 percent

Rupley

Percent of map unit: 7 percent

Morales

Percent of map unit: 5 percent

LaA—Laewest clay, 0 to 1 percent slopes

Map Unit Setting

National map unit symbol: 2v3c6

Elevation: 10 to 160 feet

Mean annual precipitation: 37 to 49 inches

Mean annual air temperature: 69 to 71 degrees F

Frost-free period: 275 to 300 days

Farmland classification: All areas are prime farmland

Map Unit Composition

Laewest and similar soils: 90 percent

Minor components: 10 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Laewest

Setting

Landform: Flats

Landform position (three-dimensional): Talf

Microfeatures of landform position: Gilgai

Down-slope shape: Linear

Across-slope shape: Linear

Parent material: Clayey fluviomarine deposits derived from igneous, metamorphic and sedimentary rock

Typical profile

A - 0 to 17 inches: clay

Bss1 - 17 to 40 inches: clay

Bss2 - 40 to 59 inches: clay

Bkss - 59 to 80 inches: clay

Properties and qualities

Slope: 0 to 1 percent

Depth to restrictive feature: More than 80 inches

Natural drainage class: Moderately well drained

Runoff class: High

Capacity of the most limiting layer to transmit water (Ksat): Very low to moderately low (0.00 to 0.06 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Calcium carbonate, maximum in profile: 41 percent

Gypsum, maximum in profile: 1 percent

Salinity, maximum in profile: Nonsaline to very slightly saline (0.0 to 3.0 mmhos/cm)

Sodium adsorption ratio, maximum in profile: 12.0

Available water storage in profile: High (about 10.8 inches)

Interpretive groups

Land capability classification (irrigated): 2s

Land capability classification (nonirrigated): 2s

Hydrologic Soil Group: D

Ecological site: Blackland 24-44" PZ (R150AY526TX)

Hydric soil rating: No

Minor Components

Dacosta

Percent of map unit: 6 percent

Landform: Flats

Landform position (three-dimensional): Talf

Down-slope shape: Linear

Across-slope shape: Linear

Ecological site: Blackland 24-44" PZ (R150AY526TX)

Hydric soil rating: No

Contee

Percent of map unit: 3 percent

Landform: Flats

Landform position (three-dimensional): Talf

Custom Soil Resource Report

Microfeatures of landform position: Gilgai
Down-slope shape: Linear
Across-slope shape: Convex
Ecological site: Blackland 24-44" PZ (R150AY526TX)
Hydric soil rating: No

Unnamed, hydric

Percent of map unit: 1 percent
Landform: Depressions
Landform position (three-dimensional): Dip
Down-slope shape: Concave
Across-slope shape: Concave
Hydric soil rating: Yes

MaC—Marcado sandy clay loam, 3 to 8 percent slopes

Map Unit Setting

National map unit symbol: dlp6
Elevation: 30 to 150 feet
Mean annual precipitation: 38 to 44 inches
Mean annual air temperature: 70 to 72 degrees F
Frost-free period: 260 to 290 days
Farmland classification: Not prime farmland

Map Unit Composition

Marcado and similar soils: 80 percent
Minor components: 20 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Marcado

Setting

Landform: Flats
Landform position (three-dimensional): Rise
Down-slope shape: Linear
Across-slope shape: Convex
Parent material: Loamy fluviomarine deposits of pleistocene age

Typical profile

H1 - 0 to 8 inches: sandy clay loam
H2 - 8 to 23 inches: clay
H3 - 23 to 80 inches: clay

Properties and qualities

Slope: 3 to 8 percent
Depth to restrictive feature: More than 80 inches
Natural drainage class: Well drained
Runoff class: Very high
Capacity of the most limiting layer to transmit water (Ksat): Very low to moderately low (0.00 to 0.06 in/hr)
Depth to water table: More than 80 inches

Custom Soil Resource Report

Frequency of flooding: None
Frequency of ponding: None
Calcium carbonate, maximum in profile: 20 percent
Gypsum, maximum in profile: 15 percent
Salinity, maximum in profile: Nonsaline to slightly saline (1.0 to 4.0 mmhos/cm)
Available water storage in profile: Moderate (about 7.1 inches)

Interpretive groups

Land capability classification (irrigated): 4e
Land capability classification (nonirrigated): 4e
Hydrologic Soil Group: D
Ecological site: Blackland 24-44" PZ (R150AY526TX)
Hydric soil rating: No

Minor Components

Dacosta

Percent of map unit: 5 percent

Laewest

Percent of map unit: 5 percent

Inez

Percent of map unit: 5 percent

Milby

Percent of map unit: 5 percent

MbB—Milby sand, 0 to 2 percent slopes

Map Unit Setting

National map unit symbol: 2vv34
Elevation: 20 to 200 feet
Mean annual precipitation: 43 to 55 inches
Mean annual air temperature: 70 to 71 degrees F
Frost-free period: 270 to 300 days
Farmland classification: Not prime farmland

Map Unit Composition

Milby and similar soils: 90 percent
Minor components: 10 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Milby

Setting

Landform: Terraces
Landform position (three-dimensional): Tread
Down-slope shape: Linear
Across-slope shape: Convex
Parent material: Sandy alluvium derived from igneous, metamorphic and sedimentary rock

Custom Soil Resource Report

Typical profile

A - 0 to 11 inches: sand
E - 11 to 28 inches: sand
Bt - 28 to 80 inches: sandy clay loam

Properties and qualities

Slope: 0 to 2 percent
Depth to restrictive feature: More than 80 inches
Natural drainage class: Moderately well drained
Runoff class: Very low
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 1.98 in/hr)
Depth to water table: About 18 to 30 inches
Frequency of flooding: None
Frequency of ponding: None
Salinity, maximum in profile: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)
Available water storage in profile: Moderate (about 6.1 inches)

Interpretive groups

Land capability classification (irrigated): 3e
Land capability classification (nonirrigated): 3e
Hydrologic Soil Group: A
Ecological site: SANDY 25-35" PZ (R150AY644TX)
Hydric soil rating: No

Minor Components

Kuy

Percent of map unit: 10 percent
Landform: Terraces
Landform position (three-dimensional): Tread
Down-slope shape: Convex
Across-slope shape: Convex
Ecological site: Deep Sand 35-42" PZ (R150AY532TX)
Hydric soil rating: No

MrA—Morales-Cieno complex, 0 to 1 percent slopes, frequently ponded

Map Unit Setting

National map unit symbol: 2v3cp
Elevation: 90 to 230 feet
Mean annual precipitation: 41 to 45 inches
Mean annual air temperature: 69 to 71 degrees F
Frost-free period: 280 to 305 days
Farmland classification: Farmland of statewide importance

Map Unit Composition

Morales and similar soils: 65 percent
Cieno and similar soils: 20 percent

Custom Soil Resource Report

Minor components: 15 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Morales

Setting

Landform: Flats

Landform position (three-dimensional): Talf, flat

Down-slope shape: Linear

Across-slope shape: Linear

Parent material: Early pleistocene loamy fluviomarine deposits derived from sedimentary rock

Typical profile

A - 0 to 4 inches: loamy fine sand

E - 4 to 8 inches: fine sandy loam

E/Bt - 8 to 15 inches: fine sandy loam

Bt/E - 15 to 19 inches: fine sandy loam

Bt1 - 19 to 29 inches: sandy clay loam

Bt2 - 29 to 63 inches: sandy clay loam

BCt - 63 to 80 inches: sandy clay loam

Properties and qualities

Slope: 0 to 1 percent

Depth to restrictive feature: More than 80 inches

Natural drainage class: Moderately well drained

Runoff class: Medium

Capacity of the most limiting layer to transmit water (Ksat): Moderately low to moderately high (0.06 to 0.20 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Salinity, maximum in profile: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)

Sodium adsorption ratio, maximum in profile: 15.0

Available water storage in profile: Moderate (about 7.9 inches)

Interpretive groups

Land capability classification (irrigated): 2w

Land capability classification (nonirrigated): 2w

Hydrologic Soil Group: C

Ecological site: Sandy Loam 28-44" PZ (R150AY542TX)

Hydric soil rating: No

Description of Cieno

Setting

Landform: — error in exists on —

Landform position (three-dimensional): Dip

Down-slope shape: Concave

Across-slope shape: Concave

Parent material: Early pleistocene loamy fluviomarine deposits derived from sedimentary rock

Typical profile

A - 0 to 7 inches: loam

E/B - 7 to 14 inches: sandy clay loam

Custom Soil Resource Report

Btg1 - 14 to 33 inches: clay loam

Btg2 - 33 to 80 inches: sandy clay loam

Properties and qualities

Slope: 0 to 1 percent

Depth to restrictive feature: More than 80 inches

Natural drainage class: Poorly drained

Runoff class: Negligible

Capacity of the most limiting layer to transmit water (Ksat): Very low to moderately low (0.00 to 0.06 in/hr)

Depth to water table: About 0 inches

Frequency of flooding: None

Frequency of ponding: Frequent

Calcium carbonate, maximum in profile: 5 percent

Salinity, maximum in profile: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)

Sodium adsorption ratio, maximum in profile: 6.0

Available water storage in profile: Moderate (about 6.7 inches)

Interpretive groups

Land capability classification (irrigated): 3w

Land capability classification (nonirrigated): 4w

Hydrologic Soil Group: D

Ecological site: Lowland 35-56" PZ (R150AY537TX)

Hydric soil rating: Yes

Minor Components

Nada

Percent of map unit: 10 percent

Landform: Flats

Landform position (three-dimensional): Talf, flat

Down-slope shape: Linear

Across-slope shape: Linear

Ecological site: Claypan Prairie 28-44" PZ (R150AY528TX)

Hydric soil rating: No

Telf

Percent of map unit: 3 percent

Landform: Flats

Landform position (three-dimensional): Talf, rise

Down-slope shape: Linear

Across-slope shape: Linear

Ecological site: Loamy Prairie 28-44 PZ (R150AY535TX)

Hydric soil rating: No

Milby

Percent of map unit: 1 percent

Landform: Terraces

Landform position (three-dimensional): Tread

Down-slope shape: Linear

Across-slope shape: Convex

Ecological site: SANDY 25-35" PZ (R150AY644TX)

Hydric soil rating: No

Nez

Percent of map unit: 1 percent

Custom Soil Resource Report

Landform: Flats
Landform position (three-dimensional): Talf, flat
Down-slope shape: Linear
Across-slope shape: Linear
Ecological site: Sandy Loam 28-40" PZ (R087AY237TX)
Hydric soil rating: No

NcA—Nada-Cieno complex, 0 to 1 percent slopes, frequently ponded

Map Unit Setting

National map unit symbol: 2v3cq
Elevation: 80 to 280 feet
Mean annual precipitation: 39 to 45 inches
Mean annual air temperature: 69 to 71 degrees F
Frost-free period: 280 to 305 days
Farmland classification: Not prime farmland

Map Unit Composition

Nada and similar soils: 70 percent
Cieno and similar soils: 20 percent
Minor components: 10 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Nada

Setting

Landform: Flats
Landform position (three-dimensional): Talf, flat
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Loamy fluviomarine deposits derived from igneous, metamorphic and sedimentary rock

Typical profile

A - 0 to 7 inches: fine sandy loam
Bt1 - 7 to 27 inches: sandy clay loam
Bt2 - 27 to 52 inches: sandy clay loam
Bt3 - 52 to 80 inches: sandy clay loam

Properties and qualities

Slope: 0 to 1 percent
Depth to restrictive feature: More than 80 inches
Natural drainage class: Moderately well drained
Runoff class: High
Capacity of the most limiting layer to transmit water (Ksat): Very low to moderately low (0.00 to 0.06 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None

Custom Soil Resource Report

Calcium carbonate, maximum in profile: 5 percent
Salinity, maximum in profile: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum in profile: 13.0
Available water storage in profile: Moderate (about 6.8 inches)

Interpretive groups

Land capability classification (irrigated): 3s
Land capability classification (nonirrigated): 3s
Hydrologic Soil Group: C
Ecological site: Claypan Prairie 28-44" PZ (R150AY528TX)
Hydric soil rating: No

Description of Cieno

Setting

Landform: — error in exists on —
Landform position (three-dimensional): Dip
Down-slope shape: Concave
Across-slope shape: Concave
Parent material: Early pleistocene loamy fluviomarine deposits derived from sedimentary rock

Typical profile

A - 0 to 7 inches: loam
E/B - 7 to 14 inches: sandy clay loam
Btg1 - 14 to 33 inches: clay loam
Btg2 - 33 to 80 inches: sandy clay loam

Properties and qualities

Slope: 0 to 1 percent
Depth to restrictive feature: More than 80 inches
Natural drainage class: Poorly drained
Runoff class: Negligible
Capacity of the most limiting layer to transmit water (Ksat): Very low to moderately low (0.00 to 0.06 in/hr)
Depth to water table: About 0 inches
Frequency of flooding: None
Frequency of ponding: Frequent
Calcium carbonate, maximum in profile: 5 percent
Salinity, maximum in profile: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum in profile: 6.0
Available water storage in profile: Moderate (about 6.7 inches)

Interpretive groups

Land capability classification (irrigated): 6w
Land capability classification (nonirrigated): 6w
Hydrologic Soil Group: D
Ecological site: Lowland 35-56" PZ (R150AY537TX)
Hydric soil rating: Yes

Minor Components

Morales

Percent of map unit: 5 percent
Landform: Flats

Custom Soil Resource Report

Landform position (three-dimensional): Talf, flat
Down-slope shape: Linear
Across-slope shape: Linear
Ecological site: Sandy Loam 28-44" PZ (R150AY542TX)
Hydric soil rating: No

Telf

Percent of map unit: 3 percent
Landform: Flats
Landform position (three-dimensional): Talf, rise
Down-slope shape: Linear
Across-slope shape: Linear
Ecological site: Loamy Prairie 28-44 PZ (R150AY535TX)
Hydric soil rating: No

Milby

Percent of map unit: 1 percent
Landform: Terraces
Landform position (three-dimensional): Tread
Down-slope shape: Linear
Across-slope shape: Convex
Ecological site: SANDY 25-35" PZ (R150AY644TX)
Hydric soil rating: No

Nez

Percent of map unit: 1 percent
Landform: Flats
Landform position (three-dimensional): Talf, flat
Down-slope shape: Linear
Across-slope shape: Linear
Ecological site: Sandy Loam 28-44" PZ (R150AY542TX)
Hydric soil rating: No

Nv—Navidad fine sandy loam, frequently flooded

Map Unit Setting

National map unit symbol: dlpc
Elevation: 50 to 550 feet
Mean annual precipitation: 38 to 44 inches
Mean annual air temperature: 70 to 72 degrees F
Frost-free period: 260 to 290 days
Farmland classification: Not prime farmland

Map Unit Composition

Navidad and similar soils: 85 percent
Minor components: 15 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Navidad

Setting

Landform: Flood plains
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Loamy alluvium of holocene age

Typical profile

H1 - 0 to 33 inches: fine sandy loam
H2 - 33 to 59 inches: loamy fine sand
H3 - 59 to 80 inches: sandy clay loam

Properties and qualities

Slope: 0 to 1 percent
Depth to restrictive feature: More than 80 inches
Natural drainage class: Well drained
Runoff class: Negligible
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 1.98 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: Frequent
Frequency of ponding: None
Calcium carbonate, maximum in profile: 5 percent
Available water storage in profile: Moderate (about 6.8 inches)

Interpretive groups

Land capability classification (irrigated): 5w
Land capability classification (nonirrigated): 5w
Hydrologic Soil Group: B
Ecological site: Loamy Bottomland 25-44" PZ (R150AY534TX)
Hydric soil rating: No

Minor Components

Chicolete

Percent of map unit: 10 percent

Ganado

Percent of map unit: 3 percent

Zalco

Percent of map unit: 2 percent